NULYTELY PREP INSTRUCTIONS revised 05/02/2023

DR. SINDHU A. ABRAHAM

(972) 867-0019

To ensure a proper and complete examination, it is necessary for you to have an empty and properly cleansed colon. This is accomplished by strictly following these instructions. YOUR PROCEDURE MAY BE CANCELLED IF THESE INSTRUCTIONS ARE NOT FOLLOWED. IF YOU HAVE PROBLEMS WITH YOUR PREP, PLEASE CALL OUR OFFICE AT (972) 867-0019

PURCHASE AT THE PHARMACY: ***BISACODYL ARE OVER THE COUNTER TABLETS***

PURCHASE <u>DULCOLAX BISACODYL pills 5MG</u> MAKE SURE IT'S THE <u>LAXATIVE NOT STOOL SOFTENER</u>

, (1) container of Nulytely (prescription attached) and Metoclopramide or Zofran ondansetron (prescription attached)—your prescription may be sent to your pharmacy electronically. <u>YOU MAY ALSO BE ASKED TO TAKE GENERIC MIRALAX DAILY 5 DAYS PRIOR TO THIS</u>
LIQUID DIET BASED ON YOUR PREVIOUS COLONOSCOPY PREP RESULTS

TWO WEEKS PRIOR TO YOUR PROCEDURE:

Stop taking St. John's Wort (an herbal supplement)

FIVE DAYS PRIOR TO YOUR PROCEDURE: STOP VITAMIN E AND FISH OIL (OMEGA 3).

Stop taking ASPIRIN or ASPIRIN-CONTAINING PRODUCTS, non-steroidal anti-inflammatories, and blood thinners. Examples include, but are not limited to: COUMADIN, PRADAXA, ELIQUIS, XARELTO, ALEVE, MOTRIN, ADVIL, IBUPROFEN, EXCEDRIN, RELAFEN, ANAPROX, LODINE, FELDENE, ECOTRIN, EMPIRIN, MOBIC, NAPROSYN, TOLECTIN, VOLTAREN, DOLOBID, MOBIC, CELEBREX, ALKA-SELTZER or PLAVIX. If you have any questions about your medications, please call our office or your pharmacy. You may take Tylenol or Acetaminophen as needed.

ONE-DAY BEFORE YOUR PROCEDURE: take your usual medications including those for blood pressure, acid reflux, heart, etc.

- 1. Remain on a clear liquid diet the entire day. You may have clear broth, Jell-O with no fruit, lemonade without pulp, Popsicles with no fruit, coffee with no cream, tea, apple juice, Gatorade, 7-UP, Sprite, Coke, Dr. Pepper, Root Beer Etc. There are no color restrictions—you may drink red or purple liquids—but keep in mind, they will come out of your rectum red or purple!
 - * NO MILK PRODUCTS OR ORANGE JUICE. DRINK AT LEAST 64 OUNCES OF FLUID (not just water) DURING THE DAY. ALCOHOL IS NOT CONSIDERED A CLEAR LIQUID!!!!!!!!!

<u>IF YOU ARE DIABETIC & ARE ON INSULIN</u> – take ½ your usual dose of insulin. If you take oral medicines for diabetes, **DO take** those as usual. **DO NOT** consume just sugar-free liquids.

- 2. At <u>7am</u> fill your gallon jug of Nulytely with water and shake well. Chill in the refrigerator. If you choose the unflavored Nulytely, you may add 2 small containers of Crystal Light (lemonade flavor) to the mixture to improve the taste.
- 3. At <u>8am</u> take your 1st dose of Metoclopramide or Zofran ondansetron by mouth.
- 4. At <u>9am</u> take 2 Bisacodyl 5mg laxative tablets. This is a laxative and will cause diarrhea.
 - ❖ While you are having bowel movements, you may want to use alcohol free baby wipes or Tucks medicated pads to wipe instead of toilet tissue. To protect the skin outside the rectum, you may use Anusol, Desitin (zinc oxide), Boudreaux's Butt Paste, or Vitamin A & D ointment.
- 5. At <u>4 pm</u> start drinking an 8oz glass of Nulytely every 15 minutes, until the entire gallon is finished. DO NOT pour over ice. It may help to drink it with a straw.
 - ❖ If you feel nauseated, it may help to take a 10−15 minute break from drinking and resume by taking sips. Continue until no later than 11pm
- 6. At **2pm and 8pm** take your 2nd and 3rd doses of Metoclopramide or Zofran ondansetron by mouth.
- 7. Do not drink anything after midnight.
 - ❖ If your procedure is scheduled for anytime after 1:30pm, you may continue to have clear liquids until 7:30am the morning of your procedure. *DO NOT* CHEW GUM OR SUCK ON CANDY WHILE YOU ARE WAITING FOR YOUR PROCEDURE.

IF YOU HAVE ANY QUESTIONS ABOUT YOUR MEDICINES OR PREP, PLEASE CALL THE OFFICE (972) 867-0019, AND WE WILL ASSIST YOU

THE DAY OF YOUR PROCEDURE:

- Take your blood pressure or heart medications at your usual time with a small sip of water, unless otherwise instructed by your physician. If you take medicines for acid reflux, also take them in the morning.
- If you have asthma, allergies, emphysema, or COPD take your allergy medicine, inhalers & medicines for that condition.
- If you are diabetic DO NOT take your diabetic medications or insulin the morning of your procedure.

AFTER THE PROCEDURE:

- ❖ You will need someone to drive you home after your procedure. You will **NOT** be allowed to take a taxi or bus home.
- YOU WILL NO BE ALLOWED TO DRIVE THE REST OF THE DAY.
- Please call our office if you need information on medical transportation services if you do not have a ride available.

LOW RESIDUE DIET TO FOLLOW 3 DAYS PRIOR TO COLONOSCOPY

Food group	Foods allowed	Foods to avoid
Milk and dairy	Milk and milk products. Includes: f cow's milk f ice cream f yogurt f cheese f cream	 f fruited yogurt f any ice cream or cheese with nuts or seeds f any milk products if you are lactose intolerant
Beverages	f coffee and tea f carbonated beverages f apple juice f strained juice f bottled water f tomato juice f fruit drinks without pulp, such as fruit punch f Kool-Aid or Hi-C (without red dye) f nutritional supplements without added fiber, such as Boost/Ensure	f any beverage containing pulp or seeds, such as orange or grapefruit juice prune juice f nutritional supplements that contain fiber
Breads, cereals, and starches	f refined breads, rolls, bagels, English muffins, pita bread, biscuits, muffins, crackers, pancakes, waffles, or pastry refined cooked and cold cereals such as hominy grits, farina, cream of wheat or rice, strained oatmeal, Cheerios, Corn/Rice Chex, Cornflakes, Rice Krispies, Special K potato and sweet potato without skin refined pasta white rice	 f whole grain breads, cereals, and pasta f oatmeal f granola f any bread, cereal, cracker, or pasta made with seeds, nuts, coconut, or raw or dried fruit f corn bread f graham crackers f brown rice f wheat germ f bran f sprouted wheat f wild rice f barley f potato skins

Food group	Foods allowed	Foods to avoid

Fruits	 f canned or cooked fruit without skins or seeds (peaches, pears, apricots, apples) f applesauce f ripe banana f jellied cranberry sauce 	 f raw fruit (bananas are okay) f canned pineapple, oranges, grapefruit sections, mixed fruit f dried fruit f all berries, melons f whole cranberry sauce f avocado f coconut
Vegetables	 f tender, well cooked fresh, canned, and frozen vegetables without seeds such as peeled carrots, green beans, and beets f strained vegetable juice f strained tomato sauce 	f all raw vegetables, such as lettuce, onion, celery, cucumber, mushrooms, scallions, etc. f vegetables with seeds f tough, fibrous cooked vegetables such as: artichokes asparagus broad beans broccoli and cauliflower brussel sprouts celery corn cucumber eggplant mushrooms onion peas (green peas) sauerkraut and cabbage spinach tomatoes zucchini summer squash, winter squash
Meat and meat substitutes	 f cooked, tender fish, poultry, beef, lamb, pork, ham, veal, organ meats f eggs f cheese f tofu f tuna fish f smooth peanut butter and other smooth nut butters 	f non-tender meats f gristle f hot dogs f salami, cold cuts f meat substitutes made with whole grains, nuts, or seeds f dried beans, peas, lentils f crunchy style peanut butter and other crunchy nut butters
Miscellaneous	 f salt, sugar, ground or flaked herbs and spices f vinegar f ketchup and mustard f soy sauce f jelly (but not jam or preserves) 	 f pepper f seed spices f seeds and nuts f coconut f popcorn f jams or preserves f pickles and olives